



OMIDRIA has separate payment status in ASCs and HOPDs for 100% of fee-for service Medicare patients*

The following information is provided to help integrate OMIDRIA into your operating routine.

OMIDRIA dilution and storage¹

Prior to dilution:

Store at 20°C to 25°C (68°F to 77°F)



Protect from light

Easy to integrate into routine operating procedures:



Add preoperatively to irrigation solution

One 4-mL single-patient-use vial to 500-mL ophthalmic irrigating solution

Can be added to irrigation solution in the surgical suite

No other preparation required

After dilution, product can be stored:



Up to 4 hours at room temperature



Up to 24 hours if refrigerated

Do not use if the solution is cloudy or contains particulate matter

*OMIDIRA qualifies for separate payment across both ASC and HOPD settings as it fulfills the requirements established in the non-opioid as a surgical supply provision by CMS https://www.federalregister.gov/documents/2024/11/27/2024-25521/medicare-and-medicaid-programs-hospital-outpatient-prospective-payment-and-ambulatory-surgical

Indications and usage

OMIDRIA® is added to ophthalmic irrigating solution used during cataract surgery or intraocular lens replacement and is indicated for maintaining pupil size by preventing intraoperative miosis and reducing postoperative ocular pain.

Important safety information

OMIDRIA must be added to irrigating solution prior to intraocular use.

OMIDRIA is contraindicated in patients with a known hypersensitivity to any of its ingredients.

Systemic exposure to phenylephrine may cause elevations in blood pressure.

Use OMIDRIA with caution in individuals who have previously exhibited sensitivities to acetylsalicylic acid, phenylacetic acid derivatives, and other nonsteroidal anti-inflammatory drugs (NSAIDs), or have a past medical history of asthma.

The most commonly reported adverse reactions at $\ge 2\%$ are eye irritation, posterior capsule opacification, increased intraocular pressure, and anterior chamber inflammation.

Please see the Full Prescribing Information for OMIDRIA at www.omidria.com/prescribinginformation.

You are encouraged to report Suspected Adverse Reactions to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.





OMIDRIA is available through national wholesalers and specialty distributors

OMIDRIA is supplied in a clear, 5-mL glass, single-patient-use vial containing 4 mL of sterile solution. The smallest sellable unit is a carton, which contains four (4) single-patient-use vials.

NDC#: 82604-600-04



×

 \Box

x4

One (1) carton contains four (4) single-patient-use vials*

x30

x120

One (1) case contains thirty (30) cartons (120 total vials)

National wholesalers and specialty distributors

McKesson Specialty Distribution*

Item number:

Contact number:

5016421 1-800-482-6700

McKesson Plasma and Biologics*

Item number:

Contact number:

2856201

1-877-625-2566

Cardinal Health Specialty Distribution[†]

Item number:

Contact number:

5875778

1-855-855-0708

Besse Medical

Item number:

Contact number:

10283186

1-888-767-7123

AmerisourceBergen Specialty Distribution[‡]

Item number:

Contact number:

10283069

Contact namber.

1-800-746-6273

FFF Enterprises, Inc.

Item number:

Contact number:

OMI4060004

1-800-843-7477

*For customers of McKesson full-line wholesaler division, OMIDRIA is available for purchase through McKesson Plasma and Biologics and McKesson Specialty Distribution.

†For customers of Cardinal full-line wholesaler division, OMIDRIA is available for purchase through Cardinal Health Specialty Distribution.

‡ For customers of AmerisourceBergen Corporation full-line wholesaler division, OMIDRIA is available through AmerisourceBergen specialty distribution and can be ordered using the Passport ordering platform through AmerisourceBergen.

For more information about OMIDRIA, please call 1–877–OMIDRIA (1–877–664–3742), visit omidria.com, or scan the QR code.

References: 1. OMIDRIA [prescribing information]. Bellevue, WA: Rayner Surgical Inc. April 2023.



